arXiv:1405.0539v4 [math.AG] 15 Apr 2016

CHARACTERISTIC FOLIATION ON NON-UNIRULED SMOOTH DIVISORS ON HYPERKÄHLER MANIFOLDS

EKATERINA AMERIK, FRÉDÉRIC CAMPANA

Abstract. We prove that the characteristic foliation $\mathcal{F}$ on a non-singular divisor D in an irreducible projective hyperkähler manifold X cannot be algebraic, unless the leaves of $\mathcal{F}$ are rational curves or X is a surface. More generally, we show that if X is an arbitrary projective manifold carrying a holomorphic symplectic 2-form, and D and $\mathcal{F}$ are as above, then $\mathcal{F}$ can be algebraic with non-rational leaves only when, up to a finite étale cover, X is the product of a symplectic projective manifold Y with a symplectic surface and D is the pull-back of a curve on this surface.

When D is of general type, the fact that $\mathcal{F}$ cannot be algebraic unless X is a surface was proved by Hwang and Viehweg. The main new ingredient for our results is the observation that the canonical class of the (orbifold) base of the family of leaves is zero. This implies, in particular, the isotriviality of the family of leaves of $\mathcal{F}$.

RÉSUMÉ: Nous montrons que si le feuilletage caractéristique $\mathcal{F}$ d'un diviseur D lisse d'une variété projective complexe symplectique irréductible X est algébrique, alors ou bien X est une surface, ou bien les feuilles de $\mathcal{F}$ sont des courbes rationnelles. Lorsque D est de type général, ce résultat est dû à Hwang et Viehweg. Nous en déduisons, lorsque X est une variété projective complexe arbitraire munie d'une 2-forme symplectique holomorphe, et D, $\mathcal{F}$ comme ci-dessus, que si les feuilles X sont des courbes algébriques non-rationnelles, alors, après revêtement étale fini, X est le produit d'une surface $K3$ ou abélienne S par une variété symplectique Y, et $D = C \times Y$ pour une courbe $C \subset S$.

L'ingrédient principal nouveau de la démonstration est l'observation que la classe canonique (orbifolde) de la base de la famille des feuilles est triviale. Ceci implique, en particulier, l'isotrivialité de la famille des feuilles de $\mathcal{F}$.

1. Introduction

Let X be a projective manifold equipped with a holomorphic symplectic form σ. Let D be a smooth divisor on X. At each point of D, the restriction of σ to D has one-dimensional kernel. This gives a non-singular foliation $\mathcal{F}$ on D, called *the characteristic foliation*. We say that $\mathcal{F}$ is *algebraic* if all its leaves are compact complex curves.

If D is uniruled, the characteristic foliation $\mathcal{F}$ is always algebraic. Indeed, its leaves are the fibres of the rational quotient fibration on D

(see for example [A-V14], Section 4). On the other hand, J.-M. Hwang and E. Viehweg proved in [H-V08] that $\mathcal{F}$ cannot be algebraic when D is of general type, except for the trivial case when $dim(X) = 2$. The aim of this article is to classify the examples where $\mathcal{F}$ is algebraic and D is not uniruled.

Our main result is as follows.

Theorem 1.1. *Let X be a projective manifold with a holomorphic symplectic form σ and let D be a smooth hypersurface in X. If $\mathcal{F}$ as above is algebraic and the genus of its general leaf is $g > 0$, then the associated fibration is isotrivial and K_D is nef and abundant, with $\nu(K_D) = \kappa(D) = 1$ when $g \geq 2$ and $\nu(K_D) = \kappa(D) = 0$ when $g = 1$.*

Here ν denotes the numerical dimension and κ the Kodaira dimension. In general, $\kappa(D)$ does not exceed $\nu(K_D)$, and K_D is said to be abundant when the two dimensions coincide (by a result of Kawamata, this implies the semiampleness of K_D, so this notion is important in the minimal model program).

What we actually are going to prove is a slightly more general result. Consider a smooth projective variety D of dimension d carrying a nowhere vanishing holomorphic $(d-1)$-form ω. Such a form has one-dimensional kernel at each point and therefore defines a smooth rank-one foliation $\mathcal{F}$. Alternatively, the foliations arising in this way are those defined by the subbundles of T_D isomorphic to the anticanonical bundle of D. In this situation, we have the following

Theorem 1.2. *If $\mathcal{F}$ is algebraic, then the associated fibration $f : D \to B$ is isotrivial without multiple fibers in codimension one and the canonical class K_B is trivial.*

We refer to subsection 2.1 for the definition and discussion of the fibration associated to a smooth algebraic foliation of rank one.

When D is a divisor in a holomorphic symplectic manifold (X, σ) of dimension $d + 1 = 2n$, one recovers the first part of theorem 1.1 by taking the form $\sigma^{\wedge(n-1)}$ for ω, since the kernel of σ is then equal to that of ω; the assertions on the numerical and Kodaira dimension are deduced from theorem 1.2 in a standard way.

The next two theorems are consequences of theorem 1.1.

Theorem 1.3. *Let X, D, $\mathcal{F}$ be as in theorem 1.1, and suppose moreover that X is irreducible (that is, simply connected and with $h^{2,0}(X) = 1$). If $\mathcal{F}$ is algebraic and D is not uniruled, then $dim(X) = 2$.*

By the Bogomolov decomposition theorem, up to a finite étale covering, any compact Kähler symplectic manifold is a product of a torus and several irreducible holomorphic symplectic manifolds. Since our assumptions on D and $\mathcal{F}$ are preserved under finite étale coverings, theorem 1.3 is valid for holomorphic symplectic manifolds with $h^{2,0} = 1$

and finite fundamental group. Moreover we may consider only the case of such products in the sequel.

Remark 1.4. *The smoothness assumption is essential, as one sees by considering the Hilbert square X of an elliptic $K3$-surface $g : S \to \mathbb{P}^1$: one has a fibration $h : X \to \mathbb{P}^2 = Sym^2(\mathbb{P}^1)$. If $C \subset \mathbb{P}^2$ is the ramification conic of the natural 2-cyclic cover $(\mathbb{P}^1)^2 \to \mathbb{P}^2$, and $L \subset \mathbb{P}^2$ is a line tangent to C, then the characteristic foliation on the singular divisor $D := h^{-1}(L)$ is algebraic with $g = 1$. One obtains similar examples with $g > 1$ by considering the image of $C \times S$ in the Hilbert square of S, where S is an arbitrary $K3$ surface and $C \subset S$ is a curve.*

Theorem 1.5. *Let X, D, $\mathcal{F}$ be as in theorem 1.1. Suppose that D is non-uniruled and $\mathcal{F}$ is algebraic. Then, possibly after a finite étale covering, $X = S \times Y$, where $dim(S) = 2$, both S and Y are complex projective manifolds carrying holomorphic symplectic forms σ_S, σ_Y, and $D = C \times Y$, where $C \subset S$ is a curve.*

Remark 1.6. *The surface S from theorem 1.5 is, up to a finite cover, either $K3$ or abelian. In the first case, $\sigma = p^*\sigma_S \oplus q^*\sigma_Y$ on $TX \cong p^*TS \oplus q^*TY$ (where p, q denote the projections) by Künneth formula.*

In the second case, one still has $\sigma = p^*\sigma_S \oplus q^*\sigma_Y$ when $g > 1$. Indeed, by Künneth formula (and Bogomolov decomposition) one reduces to the case when Y is also an abelian variety, and the decomposition then follows by a straightforward linear-algebraic computation.

In contrast to these cases, when S is an abelian surface and $g = 1$, σ is not always a direct sum (see example 4.1).

The main ideas of the proof of Theorem 1.2 are as follows. Suppose that D is not uniruled and that $\mathcal{F}$ is algebraic. Then $\mathcal{F}$ defines a holomorphic fibration $f : D \to B$ such that its non-singular fibers are curves of genus $g > 0$, and the singular fibers are multiple curves with smooth reduction. The base has only quotient singularities by Reeb stability. We prove that the codimension of the locus of multiple fibers in D (and of its image in B) is at least two. Therefore the form ω descends to B outside of a codimension-two locus; this trivializes the canonical class of B.

The generic semi-positivity theorem of [C-P13], in the simpler case when there is no orbifold structure, now implies that the Iitaka dimension of the determinant of any subsheaf of the cotangent sheaf of B is non-positive. On the other hand, Hwang and Viehweg construct such a subsheaf (coming from the Kodaira-Spencer map) with Iitaka dimension equal to the number of moduli of the fibres of f. Therefore the family f must be isotrivial.

As an application, we deduce in section 5 a certain case of the Lagrangian conjecture on a projective (and, more generally, compact Kähler) irreducible holomorphic symplectic manifold of dimension $2n$

from the Abundance conjecture in dimension $2n-1$. We therefore solve this case unconditionally for $n = 2$, since the Abundance conjecture is known for threefolds ([K92]; see also [C-H-P14] for the generalization to the Kähler case). This was our initial motivation for this research. When the research has been completed, Chenyang Xu has informed us that for projective manifolds, this case of the Lagrangian conjecture follows from a fundamental result of Demailly, Hacon and Paun ([D-H-P12]). As no algebraic proof of [D-H-P12] is known, our result also gives a simple algebro-geometric alternative for hyperkähler manifolds (see section 5 for statements and proofs).

The next section is devoted to the proof of theorems 1.1 and 1.2. The following two prove theorems 1.3 and 1.5 respectively. In the last section, we treat our application to the Lagrangian conjecture.

2. Some numerical invariants of the characteristic foliation

2.1. **Smooth rank 1 foliations.** Let D be a d-dimensional ($d \geq 2$) connected Kähler manifold carrying a non-singular holomorphic foliation $\mathcal{F}$ of rank 1. The foliation $\mathcal{F}$ is called algebraic when all its leaves are compact complex curves. A non-singular algebraic foliation induces a proper holomorphic map $f : D \to \mathcal{C}(D)$ to a component $\mathcal{C}(D)$ of the cycle space of D. Indeed, the general leaves of $\mathcal{F}$ are smooth curves varying in a dominating family of cycles on D; by compactness of $\mathcal{C}(D)$, one has well-defined limit cycles which must be supported on the special leaves, and the multiplicity of such a cycle is uniquely determined by pairing with the Kähler class. Taking the normalization of the image if necessary, we obtain a proper holomorphic map $f : D \to B$ onto a $(d-1)$-dimensional normal base B. It is well-known that in such a situation, the holonomy groups of the leaves are finite (this amounts to the boundedness of the volume of the leaves which holds in the Kähler case, see for example [E76]). Therefore by Reeb stability (see [MM], or else [H-V08] which develops the construction of [MM] in the holomorphic case in some detail), locally in some saturated neighbourhood of each leaf C of holonomy group G_C, our foliation is the quotient of $T \times \tilde{C}$, where T is a local transverse and $\tilde{C}$ is G_C-covering of C, by the natural action of G_C.

In particular, B has only quotient singularities and so is $\mathbb{Q}$-factorial, and f is "uni-smooth", that is, the reduction of any of its fibres is a smooth projective curve.

Let g denote the genus of a non-singular fiber of f. If $g = 0$, the holonomy groups are trivial and all fibres of f are smooth reduced rational curves, B is smooth, f submersive. If $g > 0$, f may have multiple fibres, of genus one when $g = 1$ and of genus greater than one (but possibly smaller than g) when $g > 1$.

If $g = 1$ and B is compact, it is well-known that f must be **isotrivial**: indeed, the j-function then holomorphically maps B to $\mathbb{C}$. In fact the holomorphicity of j near the multiple fibres is easily checked: from Reeb stability we obtain the local boundedness of j, and then use the normality of B.

A pair $D, \mathcal{F}$ as above arises, for example, when D is a smooth connected divisor in a $2n$-dimensional projective (or compact Kähler) manifold X carrying a holomorphic symplectic 2-form σ. The foliation $\mathcal{F}$ is then given, at each $x \in D$, as the σ-orthogonal to TD_x at x. In this case $d = 2n - 1$. In general, $\mathcal{F}$ will not be algebraic. One particular case when $\mathcal{F}$ is algebraic is that of a uniruled D: the leaves of $\mathcal{F}$ are then precisely the fibres of the rational quotient fibration of D (see for instance [A-V14], section 4), so $g = 0$.

We will elucidate below the situation when $\mathcal{F}$ is algebraic and $g > 0$.

Note that in this example, the quotient bundle $T_D/\mathcal{F}$ carries a symplectic form, so it has trivial determinant. Therefore the line bundle $\mathcal{F}$ is isomorphic to the anticanonical bundle of D (by adjunction, this is $\mathcal{O}_D(-D)$).

The purpose of this section is to prove the following result, which is stated as theorem 1.1 in the introduction.

Theorem 2.1. *Let D be a smooth divisor in a projective holomorphic symplectic variety (X, σ), and $\mathcal{F}$ the foliation on D given by the kernel of $\sigma|_D$. If D is non-uniruled and $\mathcal{F}$ is algebraic, then the corresponding fibration $f : D \to B$ is isotrivial, K_D is nef and abundant, $\nu(K_D) = \kappa(D) = 1$ if $g \geq 2$, and $\nu(K_D) = \kappa(D) = 0$ if $g = 1$.*

This shall be a consequence of a more general isotriviality result stated as theorem 1.2:

Theorem 2.2. *Let D be a complex projective manifold of dimension d carrying a nowhere vanishing holomorphic $(d-1)$-form ω. Let $\mathcal{F}$ be the foliation defined as the kernel of ω. Suppose $\mathcal{F}$ is algebraic. Then the corresponding fibration $f : D \to B$ is isotrivial and submersive in codimension two, and the canonical class of B is trivial.*

Our first idea is to introduce the *orbifold base* of a fibration and to show that the orbifold structure is actually trivial when the fibration is defined by a non-vanishing holomorphic $(d-1)$-form.

2.2. **Orbifold base.** Let $f : D \to B$ be a uni-smooth fibration in curves, with D smooth Kähler and B $\mathbb{Q}$-factorial. We define (as in [Ca 04], in a much more general situation there) the *orbifold base* (B, Δ) for f as follows: for each irreducible reduced Weil ($\mathbb{Q}$-Cartier) divisor $E \subset B$, set $E' = f^{-1}(E)$. This is an irreducible divisor, and $f^*(E) = m_f(E)E'$ for some positive integer $m_f(E)$. This integer is equal to 1 for all but finitely many E. Set $\Delta = \sum_{E \subset B}(1 - \frac{1}{m_f(E)})E$. The divisor Δ thus carries the information about the multiple fibers of

f in codimension one, but the coefficients of Δ are "orbifold multiplicities" varying between zero and one rather than the multiplicities of the fibers. Over a neighbourhood of a general point $b \in \Delta$ - that is, a point outside of $Sing(B)$ and $Sing(\Delta)$ - the map f is locally given by $(z_1, \dots, z_{d-1}, w) \mapsto (z_1^m, \dots z_{d-1}) = (u_1, \dots, u_{d-1})$, where $m = m_f(E)$ for the component E of Δ which contains b.

Lemma 2.3. *Suppose that f is given by the kernel of a non-vanishing holomorphic $(d-1)$-form ω. Then f has no multiple fibers in codimension one, that is, $\Delta = 0$, and K_B is trivial.*

Proof. The question on multiple fibers is local on B, and B is smooth in codimension one. We can thus assume that B is a polydisc in $\mathbb{C}^{d-1}$, with coordinates $(u_1, \dots, u_{d-1}) = (u, u')$ (u being the first coordinate and u' the $(d-2)$-tuple of others), and that f has multiple fibres of multiplicity $m > 1$ over the divisor E defined by the equation $u = 0$. Since the form ω is d-closed, and its kernel is $Ker(df)^{sat}$, the saturation being taken in TD, it descends over $B - E$ to a holomorphic $(d-1)$-form α on $B - E$ such that $\omega = f^*(\alpha)$ on $f^{-1}(B - E)$ (see e.g. [S09], lemma 6, where the full argument is given for holomorphic symplectic forms; it immediately generalizes to our setting).

We are going to show that α extends holomorphically to B, and that $m = 1$. Write $\alpha = G(u, u')du \wedge du'$ (where du' stands for the wedge product of du_i for $i > 1$). We claim that $|G(u, u')| = e^g.|u|^{-c}$, with $c = 1 - \frac{1}{m}$, where g is a real-valued bounded function, after possibly shrinking B near $(0, 0)$.

Let, indeed, $B' \subset D$ be a smooth local multisection of degree m over B meeting transversally the reduction of the fibre of f over $(0, 0) \in B$. We can choose the coordinates $(z_1, z_2, \dots z_{d-1}, w) = (z, z', w)$ on D near the intersection point $(0, 0, 0)$ of B' and the fibre $D_{(0,0)}$ of f over $(0, 0)$ in such a way that $f(z, z', w) = (z^m, z')$, and B' is defined by the equation $y = 0$. Restricting ω to B', we see that $f^*(\alpha) = G(z^m, z').m.z^{m-1}.dz \wedge dz' = \omega|_{B'} = h(z, z')dz \wedge dz'$, for some nowhere vanishing function $h(z, z') = H(z^m, z') = H(u, u')$, whenever $u = z^m \neq 0$.

Thus $|G(u, u')| = |G(z^m, z')| = \frac{|H(u,u')|}{m} . \frac{1}{|u|^c} = e^{g(u,u')} . \frac{1}{|u|^c}$.

The following well-known fact now shows that α extends holomorphically to B, and hence c must be zero and $m = 1$ as claimed.

Let $G(u, u')$ be a holomorphic function defined on $B - E$, where B is a polydisc centered at $(0, 0)$ in $\mathbb{C}^{n-1}$, and E is the divisor defined by $u = 0$ in B. Assume that, for some $\varepsilon > 0$, $|G(u, u')| \leq C.|u|^{-(1-\varepsilon)}$ for some positive constant C independent on u'. Then $G(u, u')$ extends holomorphically across the divisor $u = 0$.

Indeed, fix u', the Laurent expansion $G(u, u') = \sum_{k=-\infty}^{k=+\infty} a_k(u').u^k$ of G has then coefficients $a_k(u')r^k = \frac{1}{2\pi} \int_0^{2\pi} e^{-ikt} G(re^{it}, u')dt$ (cf. Henri Cartan, Théorie élémentaire des fonctions analytiques, Hermann 1961, p. 86, formula (2.1)). The bound on $|G|$ implies that $|a_k(u')| \leq$

$Cr^{-k-1+\varepsilon}$ for $0 < r << 1$. This implies that $a_k(u') = 0$ if $k < 0$, by letting $r \to 0^+$.

It remains to show that the Weil ($\mathbb{Q}$-Cartier) divisor K_B is trivial. Indeed the form ω descends to a non-vanishing holomorphic form on the complement of a codimension-two subset of B. Hence the triviality of K_B. □

Remark 2.4. *Another way to see this is by Reeb stability. Indeed in a neighbourhood of a multiple fiber C over a general point of Δ, which has cyclic holonomy of order m, ω must lift as a G_C-invariant form to the G_C-covering coming from Reeb stability, but this is impossible by the explicit local computation.*

Remark 2.5. *The map $f : D \to B$ given by a global nonvanishing $(d-1)$-form may have multiple fibers in codimension two: take for instance $D = (E \times E \times C)/G$ where E is an elliptic curve, C is a curve equipped with a fixed-point-free involution and G a group of order two where the non-trivial element acts as $-Id$ on $E \times E$ and as that involution on C. Then the projection onto the quotient of $E \times E$ by $-Id$ has isolated multiple fibers, and is given by the kernel of a 2-form which is the exterior product of 1-forms on E.*

2.3. **Isotriviality of the fibration.** As we have already remarked, the isotriviality of the family of curves $f : D \to B$ associated to $\mathcal{F}$ is clear when $g = 0$ or $g = 1$, so we assume in this section that $g \geq 2$. All varieties are assumed to be projective (or quasiprojective, when we work outside of a suitable codimension two subset such as $Sing(B)$). Define the sheaf Ω^1_B as the direct image $j_*\Omega^1_{B^{sm}}$ where $j : B^{sm} \to B$ is the embedding of the smooth part of B in B. The following theorem is a direct consequence of the strenghtening of Miyaoka's generic semi-positivity theorem ([Mi85], see [PM] p. 66-67, Theorem 2.14, 2.15 for a formulation adapted to our purposes) given in lemma 2.7 below.

Theorem 2.6. *Let B be a normal projective variety with log-canonical singularities such that $K_B \equiv 0$. Let L be a coherent rank-one subsheaf of $(\Omega^1_B)^{\otimes k}$ for some $k > 0$. Then $deg_C(L_{|C}) \leq 0$ for a sufficiently general complete intersection curve C cut out on B by members of a linear system $|lH|$, $l >> 0$, where H is an ample line bundle on B.*

In particular, for any integer $m > 0$ one has: $h^0(B^{sm}, L^{\otimes m}) \leq 1$ and so $\kappa(B^{sm}, det(\mathcal{F})) \leq 0$, for any coherent subsheaf $\mathcal{F} \subset \Omega^1_B$.

Proof. By lemma 2.7 below, the quotient $Q := ((\Omega^1_B)^{\otimes k}/L)$ restricted to C has non-negative degree. Thus the degree of the (locally free) sheaf $L|_C$ is non-positive, since $deg_C((\Omega^1_B)^{\otimes k}) = 0$. □

The following lemma is a special case of the main result of [C-P13] when $\Delta = 0$. Its proof is a considerably simplified version of the general case, in particular, no use of orbifold differentials is required. We refer

to [C-P13] for details (see [C-P15] for a more general result; the reader may also consult [Cl15] which exposes the main ideas and techniques of both [C-P13] and [C-P15]).

Lemma 2.7. *Let B be an n-dimensional, normal, connected, projective variety with log-canonical singularities. Assume that K_B is pseudo-effective. Then, for any $m > 0$, any quotient Q of $(\Omega^1_B)^{\otimes m}$ has non-negative slope with respect to any ample polarisation $\alpha := H^{n-1}$ of B.*

Proof. By general properties of slopes (see e.g. pages 9-10 of [Cl15]), it suffices to show that the minimal slope $\mu^{min}_\alpha(\Omega^1_B) \geq 0$ (by the *minimal slope*, one means the smallest possible slope of a quotient sheaf with respect to α); recall that we are interested in degrees of restrictions to curves C not passing through the singularities of B and therefore all sheaves we consider are locally free on such C. Assume there is a quotient of Ω^1_B with α-negative slope, then the maximal destabilising subsheaf for the dual defines an α-semistable foliation $\mathcal{F}$ on B with α-positive slope (integrability follows from Miyaoka's slope argument). By [B-MQ-01], $\mathcal{F}$ is algebraic. So there exists a rational fibration $g : B \dashrightarrow Z$ such that $\mathcal{F} = Ker(dg)$.

Taking a neat model (see [C-P13], p. 848) of g, obtained by blowing up B and Z, we get $g' : B' \to Z'$ with B', Z' smooth. We can write $K_{B'} + \Delta' = b^*(K_B) + E$, where $b : B' \to B$ is our blow-up, for Δ' and E some effective b-exceptional $\mathbb{Q}$-divisors on B' without common components. The divisor Δ' on B' is an orbifold divisor (that is has coefficients between 0 and 1) because B is log-canonical, and $K_{B'} + \Delta'$ is pseudo-effective, since so is K_B. Let $D(g', 0)$ denote the ramification divisor of g', that is, the sum of the $(g'^*F - (g'^*F)_{red})^{surj}$ over all prime divisors F of Z', where the superscript means that we consider only the non-exceptional components of $g'^*F - (g'^*F)_{red}$, i.e. those which map surjectively to F (cf. [C-P13], p.848). Theorem 2.11 of [C-P13] shows that $M := K_{B'/Z'} + (\Delta')^{hor} - D(g', 0)$ is pseudo-effective as well (here the superscript denotes the "horizontal part", dominating the base). Thus, denoting by C' the strict transform of a curve C which is a generic complete intersection of large multiples of H, we obtain $MC' \geq 0$. On the other hand, it follows from Proposition 1.9 of [C-P13] that up to a positive normalisation constant k, we have: $-det(\mathcal{F})C = k.MC'$, a contradiction since by construction $det(\mathcal{F})C > 0$. Note that though the calculation of this proposition is made in the orbifold context we actually do not need orbifold differentials: as the generic curve C avoids the singularities, C' avoids Δ', so that only the ramification of g' contributes. □

Remark 2.8. *An example where Miyaoka's result does not apply, while lemma 2.7 does, is the following Ueno surface. Let $A = E \times E$ be the product of two copies of the elliptic curve E with complex multiplication by $i = \sqrt{-1}$. Let $S := A/\mathbb{Z}_4$, the generator acting by i simultaneously*

on both factors. Then S is a rational surface with 16 quotient singularities, not all canonical. On S there is no pair $(\mathcal{F}, H)$ consisting of a rank 1 foliation $\mathcal{F}$ and a polarisation H such that the H-slope of $\mathcal{F}$ is positive. This follows from 2.7, but can also be checked directly. Indeed, otherwise both $\mathcal{F}$ and H could be lifted to A with this same intersection property, since the quotient map $q : A \to S$ is étale over S^{sm}, and A is smooth. But such a foliation does not exist on A, since its tangent bundle is trivial. The surface S is uniruled, therefore the absence of such foliations is not directly implied by [Mi85].

We return to the proof of Theorem 2.2.

The determinant of any subsheaf of Ω^1_B restricted to B^{sm} has non-positive Kodaira dimension; this also remains true for finite coverings of B, étale over B^{sm}.

Following [H-V08], we now construct a subsheaf of Ω^1_B (or more precisely of $\Omega^1_{B'}$ where B' is such a covering) such that the Kodaira dimension of its determinant over B^{sm} is equal to the variation of moduli of our family of curves; the argument is shorter here since we have remarked that f is submersive in codimension one.

Indeed, it suffices to do so outside of a codimension-two algebraic subset in B, that is, over B^0 which is smooth and such that the restriction $f : D^0 \to B^0$ of $f : D \to B$ is a smooth family of curves. It is well-known (see e.g. [H-V08], Lemma 3.1) that, after replacing B^0 by a finite étale covering, the family $f : D^0 \to B^0$ becomes the pull-back of the universal family of curves with level N structure $g : \mathcal{C}_g^{[N]} \to M_g^{[N]}$ under a morphism $j : B^0 \to M_g^{[N]}$ for a suitable $N >> 0$.

Since D^0 is now a smooth family of curves over a smooth base B^0, one can consider the "Kodaira-Spencer map"

$$f_*(\omega^{\otimes 2}_{D^0/B^0}) \to \Omega^1_{B^0}$$

obtained by dualizing the usual Kodaira-Spencer map from T_{B^0} to $R^1 f_* T_{D^0/B^0}$ associated to the family of curves $f : D^0 \to B^0$. Let $\mathcal{H} \subset \Omega^1_{B^0}$ be its image: it is a coherent subsheaf of $\Omega^1_{B^0}$. Moreover, it is functorial in B^0, that is, its construction commutes with base change.

Proposition 2.9. *(cf.* [H-V08]*, proposition 4.4) Assume that $g \geq 2$. Then $\kappa(B^0, det(\mathcal{H})) = Var(f) = dim(Im(j))$.*

Proof. The sheaf $f_*(\omega^{\otimes 2}_{D^0/B^0})$ is the pull-back by j of $g_*(\omega^{\otimes 2}_{\mathcal{C}_g^{[N]}/M_g^{[N]}})$, and the latter is ample by [H-V08], Proposition 4.3. We conclude by [H-V08], Lemma 4.2. □

Corollary 2.10. *The fibration $f : D \to B$ is isotrivial.*

Indeed, by 2.6 we know that the Kodaira dimension of the determinant of any subsheaf of $\Omega^1_{B^{sm}}$ is non-positive, and so $\kappa(B^{sm}, det(\mathcal{H})) = Var(f) = 0$. This finishes the proof of theorem 2.2.

2.4. **A more general conjectural isotriviality statement.** The corollary 2.10 is a special case of the following more general conjectural statement, which slightly generalises [T13][1]:

Conjecture 2.11. *Let $f : X \to B$ be a proper, connected, quasi-smooth[2] fibration of quasi-projective varieties, where X is smooth and B is normal. Assume that the (reduced) fibres of f have semi-ample canonical class, and that the orbifold base (B, Δ) of f is special in the following sense (cf.* [Ca 07]*): for any $p > 0$ and any coherent rank-one subsheaf $L \subset (f^*(\Omega^p_B))^{sat}$, where the saturation takes place in Ω^p_X, one has $\kappa(X, L) < p$. Then f is isotrivial.*

We would like to remark that the special case of this conjecture when $f : X \to B$ is a family of curves and the orbifold canonical bundle of the base is trivial can be proved by an argument similar to the one just given but much more subtle, using the orbifold generic semi-positivity of [C-P13] and the full argument of [H-V08]. Since this turns out to be irrelevant for the characteristic foliation by lemma 2.3, we intend to publish this elsewhere.

2.5. **Consequences of isotriviality.**

Our goal now is to get the information on K_D once the isotriviality is established. All arguments work in the compact Kähler case. Let us first remark that the relative canonical divisor $K_{D/B}$ is well-defined as a $\mathbb{Q}$-Cartier divisor, and $K_D \equiv K_{D/B}$ since K_B is trivial.

We first make a normalized base-change to remove all multiple fibers.

Lemma 2.12. *Let D be a compact connected Kähler manifold with a smooth rank-one foliation $\mathcal{F}$ with compact leaves of genus $g \geq 1$. Let $f : D \to B$ be the associated proper fibration. Consider the normalized base-change $f_D : (D \times_B D)^\nu \to D$. Then f_D is smooth.*

Proof. By definition of a foliation, a neighbourhood of $x \in D$ is isomorphic to $U' \times F$, where F is a small open subset of the leaf through x and U' is a local transverse to the foliation. Moreover, by Reeb stability, a small neighbourhood U of $b \in B$ is U'/G where G is the holonomy group, and $D'_U = (D \times_U U')^\nu$ is smooth over U' and étale over $D_U = f^{-1}(U)$. Hence $(D \times_B D)^\nu$, which locally in a neighbourhood of x is naturally isomorphic to $(D \times_U (U' \times F))^\nu = D'_U \times F$, is smooth over D: indeed the projection to D is, locally, the composition of the smooth projection to D'_U with the natural étale projection from D'_U to D_U. □

Denote by $f' : D' \to B'$ our new smooth family (so that $B' = D$ and $D' = (D \times_B D)^\nu$) and by $s : D' \to D$ the natural projection. Notice

[1] In [T13], the conjecture is established when B is smooth and $\Delta = 0$.

[2] That is, the reduction of every fibre is smooth.

that since the normalization procedure only concerns the codimension-two locus, we have $K_{D'/B'} \equiv s^* K_{D/B}$.

It is well-known that a smooth isotrivial family of curves of genus g, after a suitable finite base change, becomes a product when $g \geq 2$, and a principal fibre bundle when $g = 1$. More precisely, we have the following lemma:

Lemma 2.13. *There exists a finite proper map $h' : B'' \to B'$ such that after base-changing f' by h', we get $f'' : D'' \to B''$ and $s' : D'' \to D'$ with the following properties: $D'' \cong F \times B''$ over B'' when $g \geq 2$, and $f'' : D'' \to B''$ is a principal fibre bundle if $g = 1$. Moreover, $K_{D''/B''}$ is nef, $\kappa(D'', K_{D''/B''}) = \nu(D'', K_{D''/B''}) = 1$ if $g \geq 2$, and $\kappa(D'', K_{D''/B''}) = \nu(D'', K_{D''/B''}) = 0$ if $g = 1$.*

Here ν denotes the numerical dimension.

Proof. The smooth isotrivial family f' is a locally trivial bundle with structure group $Aut(F)$, where F is a fiber. If $g \geq 2$, this is a finite group, so that the bundle trivializes after a finite covering $h' : B'' \to B'$. If $g = 1$, we get the principal bundle structure after a finite covering corresponding to the quotient of $Aut(F)$ by the translation subgroup. The second claim is obvious when $g \geq 2$. When $g = 1$, we remark that $K_{D''/B''}$ is dual to $f''^*(R^1 f''_*(\mathcal{O}_{D''}))$, and the latter is trivial since translations on an elliptic curve operate trivially on cohomology. □

Corollary 2.14. *Let $f : D \to B$ be as above. Then K_D is nef, $\kappa(D) = \nu(D, K_D) = 1$ if $g \geq 2$, and $\kappa(D) = \nu(D, K_D) = 0$ if $g = 1$.*

Proof. Since

$$NK_{D''/B''} \equiv Ns'^*(K_{D'/B'}) \equiv Ns'^*(s^*(K_{D/B})) \equiv N(h \circ h')^*(K_D),$$

this follows from the preceding lemma, by the preservation of nefness, numerical dimension and Kodaira-Moishezon dimension under inverse images. □

This finishes the proof of the theorem 1.1 in the projective case. Remark that when $g = 1$, this argument also proves the Kähler case, since the isotriviality, for which the projectivity assumption was needed, is then automatic. This shall be used in the proof of Corollary 5.2.

In the next section, we shall give a proof of the theorem 1.3.

3. Divisors on irreducible hyperkähler manifolds.

We suppose now that X is a projective irreducible holomorphic symplectic manifold of dimension $2n \geq 4$, $D \subset X$ is a smooth non-uniruled divisor on X and the fibres of $f : D \to B$ are curves of non-zero genus tangent to the kernel of the restriction of the holomorphic symplectic form σ to D. Recall that on the second cohomology of X there is a non-degenerate bilinear form q, the *Beauville-Bogomolov form*.

By corollary 2.14 $\nu(K_D) \leq 1 < \frac{dim(X)}{2}$. On the other hand, we have the following well-known lemma (see for instance [M99], lemma 1, keeping in mind that by Fujiki formula D^{2n} is proportional to $q(D,D)^n$ with non-zero coefficient, and that the numerical dimension $\nu(D)$ of a nef divisor D is the maximal number k such that the cycle D^k is numerically non-trivial).

Lemma 3.1. *Let D be a non-zero nef divisor on an irreducible hyperkähler manifold X. Then either $\nu(D) = dim(D)$ (if $q(D,D) > 0$), or $\nu(D) = \frac{dim(X)}{2}$ (if $q(D,D) = 0$).*

Note that $\nu(X,D) = \nu(D,K_D) + 1$, since $K_D = D|_D$. Therefore $\nu(D) \leq 2$ and the only possibility is $dim(X) = 4$, $\nu(X,D) = 2$, $\nu(D,K_D) = \kappa(D) = 1$, $g \geq 2$. This case can be excluded as follows: since $\kappa(D) = \nu(D,K_D)$, D is a good minimal model and the Iitaka fibration $\phi : D \to C$ is a regular map. Its fibers S are equivalent to D^2 as cycles on X, and therefore are lagrangian. Indeed, it follows from the definition of the Beauville-Bogomolov form σ on X that

$$\int_S \sigma\bar{\sigma} = q(D,D) = 0,$$

and this implies that the restriction of σ to S is zero. So the leaves of the characteristic foliation must be contained in the fibers of ϕ, giving the fibration of S in curves of genus at least 2. But this is impossible on S, since S is a minimal surface of Kodaira dimension zero.

This proves theorem 1.3.

4. Divisors on general projective symplectic manifolds.

The purpose of this section is to prove theorem 1.5.

Recall the setting: (X,σ) is a holomorphic symplectic projective variety, $D \subset X$ is a smooth hypersurface such that its characteristic foliation $\mathcal{F}$ is algebraic and the genus g of the leaves is strictly positive. We wish to prove that up to a finite étale covering, X is a product with a surface and D is the inverse image of a curve under projection to this surface.

By Bogomolov decomposition theorem, we may assume that X is the product of a torus T and several irreducible hyperkähler manifolds H_j with $q(H_j) = 0$ (here q denotes the irregularity $h^{1,0}$) and $h^{2,0}(H_j) = 1$.

We distinguish two cases:

First case: X is not a torus. We shall proceed by induction on the number of non-torus factors in the Bogomolov decomposition of X.

Since X is not a torus, there is an irreducible hyperkähler factor H in the Bogomolov decomposition. If $X = H$, we are done. Otherwise, write $X = H \times Y$, where Y is the product of the remaining factors.

By Künneth formula, we have $\sigma_X = \sigma_H \oplus \sigma_Y$ on $TX \cong TH \oplus TY$, since $q(H) = 0$. For $y \in Y$ general, let $D_y = D \cap (H \times \{y\})$. If this is empty, then $D = H \times D_Y$ for some divisor D_Y of Y, which is smooth with algebraic characteristic foliation. Indeed, at any point of D the σ_X-orthogonal to TD is contained in the σ_X-orthogonal to $TH \subset TD$, whereas $TH^\perp = TY$ since σ_X is a direct sum. We conclude by induction in this case.

Therefore we may suppose that D dominates Y. For $y \in Y$ generic, D_y is a smooth non-uniruled divisor on $H \times y$. At any point $(h, y) \in D$ such that $D_y \neq H \times y$ is smooth at h, we have $TD_y = TD \cap TH$. Moreover, at such a point $TH \not\subset TD$ and thus, taking the σ-orthogonals, $\mathcal{F} \not\subset TY$. We get $(TD_y)^\perp = TD^\perp \oplus TH^\perp = \mathcal{F} \oplus TY$.

Since σ is a direct sum, the σ_H-orthogonal of TD_y in TH is the projection of $\mathcal{F}$ to TH.

In other words: the characteristic foliation $\mathcal{F}_{D_y}$ of D_y inside H is the projection on TH of the characteristic foliation $\mathcal{F} \subset TX$ along D_y. The leaves of $\mathcal{F}_{D_y}$ are thus the étale p_H-projections of the leaves of $\mathcal{F}$ along D_y, and so $\mathcal{F}_{D_y}$ is algebraic, with non-uniruled leaves. From theorem 1.3, we deduce that H is a $K3$-surface, and the divisors D_y are curves of genus $g > 0$ for $y \in Y$ generic.

When D_y is singular at h, one has $TH \subset TD$ at (h, y), and therefore at such points $\mathcal{F} \subset TY$.

Fix any $h \in H$ and let C_y denote the leaf of the characteristic foliation of D through (h, y). By isotriviality, all the curves C_y are isomorphic to each other. When y varies in the fibre of D over h, we thus have a positive-dimensional family of nonconstant maps $p_H : C_y \to H$ parameterized by a compact (but possibly not connected) variety D^h, and all images pass through the point $h \in H$. After a base-change $\alpha : Z \to D^h$ (not necessarily finite, but with Z still compact) of the family of the leaves, we have a map $p : C_y \times Z \to H$ mapping a section $c \times Z$ to a point. By the rigidity lemma, all images $p_H(C_y)$ coincide when y varies in a connected component of Z; therefore there is only a finite number of curves C_y through any $h \in H$. By the same reason, such a curve (that is, the projection of a leaf of $\mathcal{F}$ to H) does not intersect its small deformations in the family of the projections of leaves. The family of such curves is thus at most a one-parameter family, and there are only finitely many of them through any given point of H.

We are thus left with two cases: either all leaves of $\mathcal{F}$ project to the same curve on H, so that $p_H(D) = C \subset H$ is a curve and we are finished; or $p_H(D) = H$. In this last case, H is covered by a one-parameter family of curves C_t, which we may suppose irreducible, such that C_t does not intersect its small deformations and there is only a finite number of C_t through a given point.

Notice also that these C_t have to coincide with the connected components of the divisors D_y and therefore the generic C_t is smooth. By adjunction formula, it is an elliptic curve and H is fibered in curves C_t.

We claim that every C_t is non-singular. Indeed, suppose that some C_t is singular at $h \in H$. It has to be a connected component of a D_y for some (h, y) on a leaf of $\mathcal{F}$ projecting to C_t. As we have remarked above, the singularity of D_y at h means that $TH \subset TD$ and therefore $\mathcal{F} \subset TY$ along a connected component of $p_H^{-1}(h)$. But such a component is of strictly positive dimension and therefore would contain a leaf of $\mathcal{F}$. So there are at least two leaves of $\mathcal{F}$ through (h, y), one projecting to C_t and another to a point, which is absurd.

Since H is a $K3$-surface, it does not admit an elliptic fibration without singular fibers by topological reasons (non-vanishing of the Euler number). This is the contradiction excluding $p_H(D) = H$, and thus establishing theorem 1.5 when X is not a torus. □

Second case: $X = T$ is a torus. We shall use Ueno's structure theorem for subvarieties of tori ([U75], Theorem 10.9).

If $g > 1$, then $\kappa(D) = 1$. By Ueno's theorem there is a subtorus K of codimension 2 such that D is the inverse image of a curve on the quotient: $D = p^{-1}(C)$, where $p : T \to S := T/K$ is the projection and $C \subset S$ is a curve of genus $g' > 1$ on the abelian surface S. The σ-orthogonal space to K gives canonically a two-dimensional linear foliation $\mathcal{F}_T$ on T, such that the intersections of its leaves with D are the leaves of $\mathcal{F}$, hence smooth compact curves which project in an étale way by p onto C.

Let us show that the leaves of $\mathcal{F}_T$ are compact. Take a leaf C of $\mathcal{F}$ through a point $x \in T$. It is contained in the leaf L of $\mathcal{F}_T$ through x. Choose a group structure on T in such a way that $x = 0$. The translate of C by any point $a \in C$ is still contained in the leaf L since L is linear; on the other hand, it is not equal to C for a outside of a finite set, since $g(C) > 1$. Since L is two-dimensional and contains a family of compact curves parameterized by a compact base, L must itself be compact.

Therefore the leaves of $\mathcal{F}_T$ are translates of an abelian surface S'. It suffices now to take a finite étale base-change from S to S' to get the desired form $T' = K \times S'$, $D' = K \times C$, σ direct sum of symplectic forms on S' and K.

If $g = 1$, then $\kappa(D) = 0$, and D is a subtorus of codimension 1 with an elliptic fibration. There thus exists an elliptic curve $C \subset T$ and a quotient $\pi : T \to R = T/C$ such that $D = \pi^{-1}(V)$, where V is a codimension 1 subtorus of the torus R. Project $\rho : R \to R/V$, and consider the composition $p : T \to S := R/V$. Then S is an abelian surface, and $C' := p(C)$ is an elliptic curve on it. Moreover, $D = p^{-1}(C)$. Let K be the kernel of p: this is a subtorus of T of codimension 2. By Poincaré reducibility, there exists an abelian surface

$S' \subset T$ such that $(S' \cap K)$ is finite. After a finite étale cover, $T' = S' \times K$, and $D' = C \times K$ is of the claimed form. □

Remark 4.1. *In this last case, σ_T is in general not the direct sum of symplectic forms on S' and K. Take for example $T = S \times A$, $D = E \times A$, for $S, A, E \subset S$ Abelian varieties of dimensions $2, (n-2), 1$ respectively, with linear coordinates (x, y) on S, $(z_1, ..., z_{n-2})$ on A, and E given by $x = 0$. Take $\sigma_S := dx \wedge dy$, σ_A arbitrary on A, and $\sigma = \sigma_S + \sigma_A + dx \wedge dz$, for any nonzero linear form z on TA.*

5. Application to the Lagrangian conjecture.

Our aim is corollary 5.2 below. First we prove the following proposition.

Proposition 5.1. *Let $D \subset X$ be a smooth hypersurface in a connected compact Kähler manifold X of dimension $2n$, carrying a holomorphic symplectic 2-form σ. Denote by $\mathcal{F}$ the characteristic foliation on D defined by σ. Assume that D admits a holomorphic fibration $\psi : D \to S$ onto an $(n-1)$-dimensional connected complex manifold S, such that its general fibre is a lagrangian subvariety of X of zero Kodaira dimension. Then*

1. The foliation $\mathcal{F}$ is ψ-vertical (ie: tangent to the fibres of ψ).

2. Either the smooth fibres of ψ are tori, and then ψ is the restriction to D of a holomorphic Lagrangian fibration ψ' on some open neighborhood of D in X; or their irregularity $q(F)$ is equal to $n-1$. In this case the Albanese map $a_F : F \to Alb(F)$ is surjective and connected, and its fibres are elliptic curves which are the leaves of $\mathcal{F}$. Moreover F has a finite étale covering which is a torus.

Proof. The first claim is obvious, since, at any generic $x \in D$, the σ-orthogonal to TD_x is included into the σ-orthogonal to TF_x (where F denotes the fibre of ψ through x), which is equal to itself since F is Lagrangian.

Since the deformations of our Lagrangian fibres F cover D, we have $q(F) = h^0(Y, \Omega^1_X) = h^0(Y, N_{Y/X}) \geq dim(D) - dim(F) = n - 1$. Note that $q(F) \leq n$, since the Albanese map of a variety with zero Kodaira dimension is surjective with connected fibres by [K81].

If $q(F) = n$, F is bimeromorphic to a torus. Since it admits an everywhere regular foliation, it must be a torus. In this case F deforms in an n-dimensional family and this gives a fibration of a neighbourhood of F in X (indeed, the normal bundle to F in X is trivial since it is isomorphic to the cotangent bundle by the lagrangian condition). Otherwise, $q(F) = n - 1$ and the fibres of the Albanese map a_F are one-dimensional. In fact these are elliptic curves by $C_{n,n-1}$ ([Vi77]), and this also implies that F has a finite étale covering which is a torus.

Finally, the leaves of $\mathcal{F}$ inside F are tangent to the fibres of a_F. Indeed, since $q = n - 1$ and F moves inside an $(n-1)$-dimensional

smooth and unobstructed family of deformations (the fibres of ψ), all deformations of F stay inside D, and the natural evaluation map $ev : H^0(F, N_{F/X}) \otimes \mathcal{O}_F \to TX|_F$ must take its values in $T_{D|F}$.

Assume the leaves of $\mathcal{F}$ are not the fibres of a_F. We can then choose a 1-form u on $Alb(F)$ such that $v = a_F^*(u)$ does not vanish on $\mathcal{F}$ at the generic point z of F. The vanishing hyperplane of v_z in TF_z is however σ-dual to a vector $t_z \in TX_z$, unique and a nonzero modulo TF_z, which corresponds to the 1-form v_z under the isomorphism $(N_F)_z \cong (\Omega^1_F)_z$ induced by σ on the Lagrangian F. Since v does not vanish on $\mathcal{F}_z$ by assumption, $t_z \notin (T_D)_z$, which contradicts the fact that all first-order infinitesimal deformations of F are contained in D. □

Corollary 5.2. *Assume that X is an irreducible hyperkähler manifold of dimension $2n$, and $D \subset X$ a smooth reduced and irreducible divisor. Assume that K_D is semi-ample. Then $\mathcal{O}_X(D)$ is semi-ample.*

Proof. If the Beauville-Bogomolov square $q(D, D)$ is positive, then D is big, X is projective and the statement follows from Kawamata base point freeness theorem. So the interesting case is when D is Beauville-Bogomolov isotropic. We have $K_D = \mathcal{O}_X(D)|_D$. If K_D is semi-ample, its Kodaira dimension is equal to $\nu(K_D) = n - 1$ (lemma 3.1) and the Iitaka fibration ψ is regular. The relative dimension of ψ is equal to n. In fact $q(D, D) = 0$ implies that ψ is lagrangian in the same way as in [M01] (using that $K_D = \mathcal{O}_D(D)$ and that a suitable positive multiple $m.F$ of the fibre F is $\psi^*(H^{n-1})$ for some very ample line bundle H on S). By proposition 5.1, we have two possibilities: either F is a torus, and then the fibration ψ extends near D, since F must deform in an n-dimensional family; or F is of Albanese dimension $n - 1$ and the characteristic foliation on D is algebraic.

In the first case we conclude by [G-L-R 11], [H-W12] and [M08]. In the second case, we notice that since F has numerically trivial canonical bundle, the fibers of the characteristic foliation, which by proposition 5.1 are tangent to F, must be elliptic curves by adjunction formula. Therefore the characteristic foliation is isotrivial, and corollary 2.14 together with the proof of theorem 1.3 imply that this is impossible unless in the case $n = 1$, which is well-known.

□

Recall that the Lagrangian conjecture affirms that a non-zero nef Beauville-Bogomolov isotropic divisor is semiample (and thus there is a lagrangian fibration associated to some multiple of such a divisor). Corollary 5.2 shows that the Lagrangian conjecture is true for an effective smooth divisor on a holomorphic symplectic manifold of dimension $2n$, if the Abundance conjecture holds in dimension $2n - 1$. Since the Abundance conjecture is known in dimension 3, we have the following:

Corollary 5.3. *Let X be an irreducible hyperkähler manifold of dimension 4, and D a nef divisor on X. Assume that D is effective and smooth. Then $\mathcal{O}_X(D)$ is semi-ample.*

Notice that if $dim(X) = 4$, we can use [A11] instead of [G-L-R 11] and [H-W12], and [AC05] instead of [M08], so that the proof becomes more elementary in this case.

Acknowledgements: We are grateful to Jorge Pereira who suggested to look at $d-1$-forms rather than at 2-forms, and to Michael McQuillan who asked a question which led us to a simplification of the original argument.

The first author's research was carried out within the National Research University Higher School of Economics Academic Fund Program for 2015-2016, research grant No. 15-01-0118.

References

[A11] E. Amerik. A remark on a question of Beauville about lagrangian fibrations. Moscow Mathematical Journal 2012, Vol. 12, No. 4, p. 701–704.

[AC05] E. Amerik, F. Campana. Fibrations méromorphes sur certaines variétés à fibré canonique trivial. Pure and Applied Mathematics Quarterly 4(2) (2008). P. 1-37.

[A-V14] E. Amerik, M. Verbitsky. Rational curves on hyperkähler manifolds. International Math. Research Notices (2015) Vol. 2015 no. 23, 13009-13045

[B-MQ-01] F. Bogomolov-M. McQuillan.Rational curves on foliated varieties. IHES preprint. IHES/M/01/07. Février 2001.

[Ca 04] F. Campana. Orbifolds, special varieties and classification theory. Ann. Inst. Fourier 54 (2004), 499-665.

[Ca 07] F. Campana. Orbifoldes géométriques spéciales et classification biméromorphe des variétés Kählériennes compactes. JIMJ 10 (2011), 809-934.

[C-H-P14] F. Campana, A. Höering, T. Peternell. Abundance for Kähler threefolds. arXiv 1403.3175.

[C-P13] F. Campana, M. Paŭn. Orbifold generic semi-positivity: an application to families of canonically polarised manifolds. arXiv 1303.3169. Ann. Inst. Fourier. 65 (2015), 835-861 .

[C-P15] F. Campana, M. Paŭn. Foliations with positive slopes and birational stability of the orbifold cotangent bundle. arXiv 1508.02456. .

[Cl15] B. Claudon, Positivité du cotangent logarithmique et conjecture de Shafarevich-Viehweg, Seminaire Bourbaki 2015-2016, exposé 1105, http://www.bourbaki.ens.fr/TEXTES/1105.pdf

[D-H-P12] J.P. Demailly, C. Hacon, M. Paŭn. Extension theorems, non-vanishing and the existence of good minimal models. Acta Math. 210 (2013), no. 2, 203259.

[E76] D. Epstein. Foliations with all leaves compact. Ann. Inst. Fourier (Grenoble) 26 (1976), no. 1, 265282.

[G-L-R 11] D.Greb, C. Lehn, S. Rollenske. Lagrangian fibrations on hyperkähler manifolds—on a question of Beauville, Ann. Sci. ENS, 46, (2013), 375–403.

[H-V08] J.M. Hwang, E. Viehweg. Characteristic foliation on a hypersurface of general type in a projective symplectic manifold, Compos. Math. 146, (2010), 497-506.

[H-W12] J.M. Hwang, R. Weiss. Webs of Lagrangian tori in projective symplectic manifolds, Invent. Math., 192, (2013), 83-109.

[K81] Y. Kawamata. Characterization of abelian varieties. Compositio Math. 43 (1981), no. 2, 253276.

[K92] Y. Kawamata. Abundance theorem for minimal threefolds. Inv. Math. 108 (1992), 229-246.

[M99] D. Matsushita. On fibre space structures of a projective irreducible symplectic manifold, Topology 38 (1999), no. 1, 7983.

[M01] D. Matsushita. Addendum: On fibre space structures of a projective irreducible symplectic manifold, Topology 40 (2001), 431-432.

[M08] D. Matsushita. On nef reduction of a projective irreducible symplectic manifold, Math. Zeit. 258, (2008), 267-270.

[Mi85] Y. Miyaoka. Deformation of a morphism along a foliation. Algebraic Geometry Bowdoin 1985. Proc. Symp. Pure Math. 46 (1987), 245–268.

[MM] I. Moerdijk, J. Mrcun: Introduction to foliations and Lie groupoids, Cambridge University Press 2003.

[PM] Y. Miyaoka, Th. Peternell. Geometry of higher-dimensional algebraic varieties. DMV Seminar, 26, Birkhuser Verlag, Basel, 1997.

[S09] J. Sawon. Foliations on hypersurfaces in holomorphic symplectic manifolds, Int. Math. Res. Not. IMRN 23, (2009), 4496-4545.

[T13] B. Taji. The isotriviality of families of canonically-polarised manifolds over a special quasi-projective base. arXiv 1310.5391

[U75] K. Ueno.Classification Theory of Algebraic Varieties and Compact Complex Spaces. LNM 439 (1975), Springer Verlag.

[Vi77] E. Viehweg. Canonical divisors and the additivity of the Kodaira dimension for morphisms of relative dimension one. Comp. Math. 35(1977), 197-233.

Institut Elie Cartan, Université Henri Poincaré, B. P. 70239, F-54506 Vandoeuvre-lès-Nancy Cedex, France, and: Institut Universitaire de France

E-mail address: frederic.campana@univ-lorraine.fr

National Research University Higher School of Economics, Department of Mathematics, Laboratory of Algebraic Geometry, Vavilova 7, 117332 Moscow, Russia

E-mail address: ekaterina.amerik@gmail.com